



BCSortho.com

DRCL

POSTOPERATIVE INSTRUCTION SHEET – SHOULDER Replacement

Patient name _____

Procedure: _____

Sling Instructions: Patient is to follow the instructions for # checked.
A sling or shoulder immobilizer has been placed on your shoulder to minimize discomfort and protect any rotator cuff or labral repair. To avoid developing stiffness remove your arm from the immobilizer 4 to 5 times per day to straighten your elbow, and move your wrist, and hand ONLY.

- #1** Shoulder immobilizer may be removed for showering. Keep arm in the same position as if it were in the immobilizer or let the arm hang straight down when showering. Wear sling at all other times including sleeping.

Ice (cryotherapy)

Icing is most important in the first 72 hours to reduce postoperative pain and inflammation in the operative area. We recommend continuing to use ice machine liberally the first week, even at night for 20-30 minute time periods. During the 2nd and 3rd week use the ice machine 2 to 3 times a day for 20-30 minutes will help lessen the pain/inflammation.

How should I manage my surgical site, dressing and steri-strips?

- **You have a special dressing that contains silver which is a natural antimicrobial. Leave your dressing on until your follow up appointment**
- **You may shower with the dressing in place (No submerging the shoulder in water however, i.e. hot tubs, bath tubs).**
- **If the bandage becomes saturated underneath, remove the dressing and call the office.**



Activities:

Many shoulder patients find that lying down increases their discomfort. You may find that it is easier to sleep in a recliner or propped up in bed with a pillow supporting your elbow in a comfortable position for a few days. A sling or shoulder immobilizer has been placed on your shoulder to minimize discomfort and protect the repaired muscles. To avoid developing stiffness, remove your arm from the immobilizer 4 to 5 times per day to straighten your elbow, and move your wrist, and hand.

- a) Let pain be your guide to activity. (too much pain = too much activity)
- b) DO NOT use exercise machines unless specified.
- c) Squeeze a small ball as much as possible to help decrease swelling.
- d) Generally if you have a job with little physical activity, you may return to work on the third post-op day with no use of the affected arm.
- e) If your job requires excessive lifting or use of your arm, then discuss return to work date with your physician

May I use my involved arm for dressing, bathing, driving, and other daily activities? How should I manage my arm in the shower?

- You may use your wrist, hand, and elbow for daily activities. This includes eating, shaving, dressing, as long as you do not move your operated arm away from your body and it does not increase your pain.
- Do not use your arm to push up/off the bed or chair for 8 weeks after your surgery.
- When using your keyboard and mouse, do not move your arm away from your body.
- When showering, you may wash under the involved arm pit by bending forward to let the involved arm hang freely and reaching under with the opposite arm
- Do not actively move your arm away from your body.

After a shoulder replacement one of the most important factors in your outcome is getting the muscle on the front of your shoulder to heal back to bone. You should NOT actively raise or externally rotate your arm past neutral for 8 weeks. Your physical therapist will move the shoulder for you per your specific protocol. A detailed rehabilitation protocol will be provided at the first post-operative visit specific to the type of surgery that was performed.



BCSortho.com

DRCL

You may bend and straighten your elbow to perform activities such as eating or using a computer. Do not lift anything heavier than a cup of coffee.

Pain may take days to weeks to completely subside and varies depending upon the extent of the damage that needed to be repaired. Swelling will typically decrease over the first few days to first week. If you experience severe and worsening pain, redness, drainage of something other than water or watery bloody material, or fever over 101 (after the first day or two), this could indicate infection, and you should contact Dr. Luedke's office if you have any concerns.

Can I smoke following surgery?

- You should not smoke after surgery as it interferes with tendon healing.

EMERGENCIES • If you have an emergency contact Dr. Luedke's office at (979) 776-0169 and he will be contacted.

- Contact the office if you notice any of the following:
 - Uncontrolled nausea or vomiting
 - reaction to medication
 - fever greater than 101.5 (low grade fevers 1-2 days after surgery is normal)
 - severe pain not relieved by pain medication/ice/elevation of leg
 - redness or continued drainage around incisions (a small amount is normal for a few days)
 - calf pain or severe swelling.
- **If you are having chest pain or difficulty breathing, call 911 or go to the closest emergency room.**